Google scholar tuberculin antigen adjuvant acrylate OR acrylic

Search

Advanced Scholar Search Scholar Preferences

Scholar Articles and patents

anytime

include citations

Create email alertResults 1 - 10 of about 505. (0.17 sec)

An update on vaccine adjuvants in clinical trial

R EDELMAN - AIDS research and human retroviruses, 1992 - liebertonline.com

... about adjuvant safety has encouraged the development of unique synthetic antigen constructs

(Table ... basement membranes or neurolemma, causing glomerulonephritis or

meningoencephalomyelitis • Sensitization to tuberculin or to ... UPDATE ON VACCINE ADJUVANT ...

Cited by 26 - Related articles - BL Direct - All 3 versions

Survey of human-use adjuvants

RT Kenney, R Edelman - Expert Rev Vaccines, 2003 - expert-reviews.com

... Acetylated polymannose (Acemannan), Sulfolipopolysaccharide Polymethyl methacrylate (PMMA),

Acrylic acid-allyl ... neurolemma, causing glomerulonephritis or meningoencephalitis • Sensitization

to tuberculin or to ... of the adjuvant alone and the antigen-adjuvant combina- tion ...

Cited by 44 - Related articles - All 4 versions

Adjuvant-induced selective recruitment of specifically reactive lymphocytes: a probable mechanism of adjuvant action

EE Emeson - The Journal of Immunology, 1980 - Am Assoc Immnol

... All paw injections were given in 40 pl volumes with tuberculin syringes equipped with 27 ... of the tissues of a control group challenged with a third party indifferent antigen. ... RESULTS Experiment

1. Time relationship between the administration of adjuvant and semiallogeneic cells. ...

Cited by 4 - Related articles - All 2 versions

Adjuvants

R Edelman, CO Tacket - International reviews of immunology, 1990 - informaworld.com

... Sensitization to tuberculin or to other skin test antigens Carcinogenesis Pyrogenesis

Teratogenesis Abortogenesis ... vaccinated subject, and intrinsic characteristics of the antigen. ... The carrier/adjuvant/ antigen formulation should be hand-tailored and then standardized so that it ...

Cited by 18 - Related articles - All 4 versions

Delayed-type hypersensitivity after immunization with ultrasonicated Mycobacterium lepraemurium in C3H and C57BL mice

M Løvik, OA Haugen, O Closs - Scandinavian journal of ..., 1984 - interscience.wiley.com

... Adjuvant effect of DTH to MLMSon-S on induction of DTH to an unrelated antigen DTH reactions may exert an adjuvant effect on the development of DTH to an unrelated second antigen [5], and this has been reported to distinguish tuberculin-type DTH from Jones-Mote ...

Cited by 4 - Related articles - All 4 versions

A Mr 95,000 polypeptide in Porphyridium cruentum phycobilisomes and thylakoids: Possible function in linkage of phycobilisomes to thylakoids and in energy transfer

pnas.org [PDF]
Full View

T Redlinger, E Gantt - ... of the National Academy of Sciences, 1982 - National Acad Sciences ... Booster shots (0.5 mg of **antigen** in Freund's **adjuvant**) were given after 3- and 6-week intervals. ... Rabbit preimmune serum was negative when tested against the **antigen**. ... collected in the void volume after passage through a Seph- adex G-50 column (in a 1 ml **tuberculin** syringe). ...

Cited by 56 - Related articles - All 6 versions

Polymer-metallic complexes of mycobacterial protein antigen prevent dissemination of postvaccinal infection in T-deficiency states

AS Norimov, MI Mustafaev, Y Romanova, TV ... - Bulletin of Experimental ..., 1991 - Springer ... these conditions immunization with TBP in aqueous emulsion mixture of Freund's adjuvants, both complete and incomplete, did not lead to the formation of a protective effect. Thus a triple immunogenic complex consisting of tuberculous protein antigen, CP of acrylic acid and ... Related articles - All 2 versions

lontophoretic device and method for administering immune response-enhancing agents and compositions

A Matsumura, SG Reed - US Patent App. 11/280,805, 2005 - Google Patents ... No. 5,980,898 describes a patch for passive transcutaneous immunization comprising an **antigen**, an **adjuvant** and a dressing. ... In particular, **adjuvants** are administered with vaccines or antigens to enhance the immune response to the vaccine or **antigen**. ... All 3 versions


Listeria cell wall fraction. Characterization of in vitro adjuvant activity.


nih.gov [PDF]

C Schuffler, PA Campbell - Immunology, 1976 - ncbi.nlm.nih.gov ... Briefly, 25 ml of washed Degalon V26 polymethylmeta-acrylic beads (Degussa Wolf- gang Au, Germany) were ... of 3 days than at the end of 4 days and since significant adjuvant effects were ... set up as previously described, with or without 30 pg of LCWF and SRBC as antigen. ... Cited by 7 - Related articles - All 3 versions

Multiple Emulsions: Delivery System for Antigens

A BOZKIR, OM SAKA - Multiple emulsions: technology and ..., 2008 - books.google.com ... Accidental self-inoculation can result in **tuberculin** sensitization followed by chronic local inflammation ... **adjuvant** without the inclusion of muramyl dipeptide Unlike Freund's complete **adjuvant** (FCA), which ... to work with, SAF is fluid and is easily mixed with **antigen** solutions (Hjorth ... Related articles - All 2 versions





tuberculin antigen adjuvant acrylate Search

Go to Google Home - About Google - About Google Scholar

©2010 Google